



**Section 2**  
**Effects on Biotic Systems**

**Test Guideline No. 218**  
**Sediment-Water Chironomid Toxicity**  
**Using Spiked Sediment**

4 July 2023

**OECD Guidelines for the**  
**Testing of Chemicals**



## OECD GUIDELINE FOR THE TESTING OF CHEMICALS

### Sediment-water Chironomid Toxicity Test Using Spiked Sediment

#### INTRODUCTION

1. This Test Guideline is designed to assess the effects of exposure of the sediment-dwelling larvae of the freshwater dipteran *Chironomus* sp to chemicals. It is based on existing toxicity test protocols for *Chironomus riparius* and *Chironomus dilutus* (previously named *C. tentans*) which have been developed in Europe (1)(2)(3) and in North America (4)(5)(6)(7)(8) and ring-tested (1)(6)(9). Other well documented chironomid species may also be used, for example *Chironomus yoshimatsui* (10)(11).

2. The exposure scenario used in this guideline is spiking of sediment with test chemical. The selection of the appropriate exposure scenario depends on the intended application of the test. The scenario of spiking sediment is intended to simulate accumulated levels of chemicals persisting in the sediment. This exposure system involves spiking sediment of a sediment-water test system.

3. Chemicals that need to be tested towards sediment-dwelling organisms usually persist in this compartment over long time periods. The sediment-dwelling organisms may be exposed via a number of routes (i.e. waterborne, sediment and dietary exposure). The relative importance of each exposure route, and the time taken for each to contribute to the overall toxic effects, is dependent on the physical-chemical properties of the chemical concerned. For strongly adsorbing chemicals (e.g. with log Kow > 5) or for chemicals covalently binding to sediment, ingestion of contaminated food may be a significant exposure route. In order not to underestimate the toxicity of highly lipophilic chemicals, the use of food added to the sediment before application of the test chemical is expected to be considered. In order to take all potential routes of exposure into account the focus of this Guideline is on long-term exposure. The test duration is in the range of 20 - 28 days for *C. riparius* and *C. yoshimatsui*, and 28 - 65 days for *C. dilutus*. If short-term data are required for a specific purpose, for example to investigate the effects of unstable chemical, additional replicates may be removed after a 10-day period.

4. The measured endpoints are based on the following biological variables: the number of emerged males, the number of emerged females (i.e. thus also the total number of adults emerged) as well as the time to emergence. Measurements of larval survival and growth should only be made after a 10-day period if additional short-term data are required, using additional replicates as appropriate.

5. The use of formulated sediment (also called reconstituted, artificial or synthetic sediment) is recommended. Formulated sediment has several advantages over natural sediments:

- the experimental variability is reduced because it forms a reproducible "standardised matrix" and the need to find uncontaminated and clean sediment sources is eliminated;
- the tests can be initiated at any time without encountering seasonal variability in the test sediment and there is no need to pre-treat the sediment to remove indigenous fauna; the use of formulated

sediment also reduces the cost associated with the field collection of sufficient amounts of sediment for routine testing;

- the use of formulated sediment allows for comparisons of toxicity and ranking chemicals accordingly.

6. Definitions used are given in Annex 1.

## PRINCIPLE OF THE TEST

7. First-instar chironomid larvae are exposed to a concentration range of the test chemical in sediment-water systems. The test chemical is spiked into the sediment and first-instar larvae are subsequently introduced into test beakers in which the sediment and water concentrations have been stabilised. Chironomid emergence and development rate is measured at the end of the test. Larval survival and weight may also be measured after 10 days if required (using additional replicates as appropriate). These data are analysed either by using a regression model in order to estimate the concentration that would cause x % reduction in emergence or larval survival or growth (e.g. EC<sub>10</sub>, EC<sub>50</sub> etc.), or by using statistical hypothesis testing to determine a NOEC/LOEC. The latter requires comparison of effect values with control values using statistical tests. Depending on the requirements of the respective regulatory framework, the test design should be optimised to estimate an EC<sub>x</sub> endpoint and/ or to determine a NOEC/LOEC (see section TEST DESIGN).

## INFORMATION ON THE TEST CHEMICAL

8. The water solubility of the test chemical, its vapour pressure, measured or calculated octanol-water partition coefficient and partitioning into sediment and stability in water and sediment should be known. A reliable analytical method for the quantification of the test chemical in overlying water, pore water and sediment with known and reported accuracy and limits of detection and quantification should be available. Useful information includes the structural formula and purity of the test chemical. Chemical fate of the test chemical (e.g. dissipation, abiotic and biotic degradation, etc.) also is useful information. Further guidance for testing chemicals with physical-chemical properties that make them difficult to perform the test is provided in (12).

## REFERENCE SUBSTANCES

9. The testing of a reference chemical should be carried out regularly to assure that the test protocol and test conditions are reliable. Examples of reference toxicants used successfully in ring-tests and validation studies are: lindane, trifluralin, pentachlorophenol, cadmium chloride and potassium chloride (1)(2)(5)(6)(13).

## VALIDITY OF THE TEST

10. For the test to be valid the following conditions should apply:

- the emergence in each control replicate should be at least 70 % at the end of the test (1) (6); please note that data should be reported for each replicate (see paragraph 54);

- *C. riparius* and *C. yoshimatsui* emergence to adults from control vessels should occur between 12 and 23 days after their insertion into the vessels; for *C. dilutus*, a period of 20 to 65 days is necessary;
- throughout the test period, the oxygen concentration should be at least 60 % of the air saturation value (ASV) at the temperature used, and the pH of overlying water should be between 6 and 9;
- throughout the test period, the water temperature should not differ by more than  $\pm 1.0$  °C.

## DESCRIPTION OF THE METHOD

### Test Vessels

11. The study is conducted in glass 600 ml beakers measuring 8 cm in diameter. Other vessels are suitable, but they should guarantee a suitable depth of overlying water and sediment. The sediment surface should be sufficient enough to provide 2 to 3 cm<sup>2</sup> per larvae. The ratio of the depth of the sediment layer to the depth of the overlying water should be 1:4. Test vessels and other apparatus that will come into contact with the test system should be made entirely of glass or other chemically inert material (e.g. polytetrafluoroethylene (PTFE)).

### Test Species

12. The species to be used in the test is preferably *C. riparius*. *C. dilutus* is also suitable but more difficult to handle and requires a longer test period. *C. yoshimatsui* may also be used. Details of culture methods are given in Annex 2 for *C. riparius*. Information on culture conditions is also available for other species, i.e. *C. dilutus* (4) (14) and *C. yoshimatsui* (11). Identification of species should be confirmed before testing but is not required prior to every test if organisms come from an in-house culture.

### Sediment

13. Formulated sediment should preferably be used. However, if natural sediment is used, it should be characterised (at least, pH and organic carbon content should be measured; in addition, it is recommended to determine other parameters such as C/N ratio and granulometry) and it should be free from any contamination and other organisms that might compete with, or consume the chironomids. It is also recommended that, before it is used in a chironomid toxicity test, the sediment be conditioned for seven days under the same conditions which prevail in the subsequent test.

The following formulated sediment, based on the artificial soil used in Guideline 222 (15), is recommended for use in this test (1)(16)(17):

- a) 4 - 5 % (dry weight) peat (e.g. Sphagnum): as close to pH 5.5 to 6.0 as possible; it is important to use peat in powder form, finely ground (particle size  $\leq 1$  mm) and only air dried.
- b) 20 % (dry weight) kaolin clay (kaolinite content preferably above 30 %).
- c) 75 - 76 % (dry weight) quartz sand (fine sand should predominate with more than 50 % of the particles between 50 and 200  $\mu\text{m}$ ).
- d) Deionised water is added to obtain a sediment with a water content in a range of 30-50%.
- e) Calcium carbonate of chemically pure quality ( $\text{CaCO}_3$ ) is added to adjust the pH of the final mixture of the sediment to  $7.0 \pm 0.5$ .

- f) Organic carbon content of the final mixture should be 2 % ( $\pm 0.5$  %) and is to be adjusted by the use of appropriate amounts of peat and sand, according to (a) and (c).

14. The source of peat, kaolin clay and sand should be known. The sediment components should be checked for the absence of chemical contamination (e.g. heavy metals, organochlorine compounds, organophosphorus compounds, etc.). An example for the preparation of the formulated sediment is described in Annex 3. Mixing of dry constituents is also acceptable if it is demonstrated that after addition of overlying water a separation of sediment constituents (e.g. floating of peat particles) does not occur, and that the peat or the sediment is sufficiently conditioned.

15. Sediment type and sediment properties can substantially affect the fate and bioavailability of the test chemical, and therefore uptake by the test organism and subsequent toxicity. If natural sediment is used, it is necessary that sediment characteristics that may have an influence on bioavailability and therefore toxicity are reported. Total organic carbon usually provides the main binding phase for many types of chemicals and therefore its concentration in natural sediment should be measured so that normalisation of toxicity data to organic carbon can be done (18) (19) (see paragraph 45).

## Water

16. Reconstituted water should preferably be used. However, any water which conforms to the chemical characteristics of acceptable dilution water as listed in Annexes 2 and 4 is suitable as test water. Any suitable water, natural water (surface or ground water), reconstituted water (see Annex 2) or dechlorinated tap water are acceptable as culturing water and test water if chironomids will survive in it for the duration of the culturing and testing without showing signs of stress (e.g., unusual behaviour). At the start of the test, the pH of the test water should be between 6 and 9 and the total hardness not higher than 400 mg/l as CaCO<sub>3</sub>. However, if there is an interaction suspected between hardness ions and the test chemical, lower hardness water should be used (and thus, Elendt Medium M4 should not be used in this situation). The same type of water should be used throughout the whole study. The water quality characteristics listed in Annex 4 should be measured at least twice a year if natural or tap water is used or when it is suspected that these characteristics may have changed significantly. If using natural water, special care should be taken considering the characteristics mentioned.

## Stock solutions - Spiked sediments

17. Spiked sediments of the chosen concentration are usually prepared by addition of a solution of the test chemical directly to the sediment. A stock solution of the test chemical dissolved in deionised water is mixed with the formulated sediment by rolling mill, feed mixer or hand mixing. If the test chemical is poorly soluble in water, the use of a solvent may not be avoidable, i.e. it is needed to produce a suitable concentrated stock solution. In such case, the Guidance Document No. 23 (12) should be consulted. Examples of suitable organic solvent (i.e. solvent with low toxicity) are hexane, acetone or chloroform. This solution is then mixed with 10 g of fine quartz sand for one test vessel. The solvent is allowed to evaporate and it has to be totally removed from sand; the sand is then mixed with the suitable amount of sediment per test beaker. Only agents which volatilise readily can be used to solubilise, disperse or emulsify the test chemical. It should be born in mind that the sand provided by the test chemical and sand mixture, has to be taken into account when preparing the sediment (i.e. the sediment should thus be prepared with less sand). Care should be taken to ensure that the test chemical added to sediment is thoroughly and evenly distributed within the sediment. If necessary, subsamples can be analysed to determine degree of homogeneity.

## TEST DESIGN

18. The test design relates to the selection of the number and spacing of the test concentrations, the number of vessels at each concentration and the number of larvae per vessel. Designs for EC point estimation, for estimation of NOEC, and for conducting a limit test are described.

### Design for analysis by regression

19. The effect concentration (e.g. EC<sub>10</sub>, EC<sub>50</sub>) and the concentration range, over which the effect of the test chemical is of interest, should be spanned by the concentrations included in the test. Generally, the accuracy and especially validity, with which estimates of effect concentrations (EC<sub>x</sub>) can be made, is improved when the effect concentration is within the range of concentrations tested. Extrapolation outside the concentration range tested should be avoided. A preliminary range-finding test is helpful for selecting the range of concentrations to be used (see paragraph 28).

20. If the EC<sub>x</sub> is to be estimated, at least five concentrations and three replicates for each concentration should be tested. In any case, it is advisable that sufficient test concentrations are used to allow good model estimation. The factor between concentrations should not be greater than two (an exception could be made in cases when the dose response curve has a shallow slope). The number of replicates at each treatment can be reduced in some cases; this is when the number of test concentrations with expected different responses is largely increased. Increasing the number of replicates or reducing the size of the test concentration intervals tends to lead to narrower confidence intervals for the test. Additional replicates are required if 10-day larval survival and growth are to be estimated.

### Design for estimation of a NOEC/LOEC

21. If the LOEC or NOEC are to be estimated, five test concentrations with at least four replicates should be used and the factor between concentrations should not be greater than two. The number of replicates should be sufficient to ensure adequate statistical power to detect a 20 % difference from the control at the 5 % level of significance ( $p = 0.05$ ). With the development rate, an Analysis of Variance (ANOVA) is usually appropriate, such as Dunnett-test and Williams-test (20)(21)(22)(23). For the emergence data, the Cochran-Armitage, Fisher's exact (with Bonferroni correction), or Mantel-Haentzel tests may be used. Alternatively, the CPFISH (closure principle and Fisher-Freeman-Halton) test may be used since this test has a high statistical power (24).

#### Limit test

22. A limit test may be performed (one test concentration and control) if no effects were seen in the preliminary range-finding test. The purpose of the limit test is to perform a test at a concentration sufficiently high to enable decision makers to exclude possible toxic effects of the chemical, and the limit is set at a concentration which is not expected to appear in any situation. A concentration of 1000 mg/kg dry sediment is recommended. Usually, at least six replicates for both the treatment and control are necessary. Adequate statistical power to detect a 20 % difference from the control at the 5 % level of significance ( $p = 0.05$ ) should be demonstrated. With metric response (development rate and weight), the t-test is a suitable statistical method if data meet the requirements of this test (normality, homogeneous variances). The unequal-variance t-test or a non-parametric test, such as the Wilcoxon-Mann-Whitney test may be used, if these requirements are not fulfilled. With the emergence ratio, the Fisher exact test is appropriate.

## PROCEDURE

### Conditions of exposure

#### **Preparation of spiked sediment - water system**

23. The spiked sediments are placed in the vessels and overlying water is added to produce a sediment-water volume ratio of 1:4 (see paragraphs 11 and 15). The depth of the sediment layer should be of  $1.5 \pm 0.25$  cm. The weight of the sediment added to each test chamber (or at least the test chambers being used for analytical confirmation) should be recorded. This information is needed to calculate the mass balance. To avoid separation of sediment ingredients and re-suspension of fine material during addition of test water in the water column, the sediment can be covered with a disc (made of e.g. stainless steel or PTFE) while water is poured onto it, and the disc removed immediately afterwards. Other devices may also be appropriate.

24. The test vessels should be covered (e.g. glass plates with openings for air tubing). If necessary, during the study the water levels will be topped to the original volume in order to compensate for water evaporation. This should be performed using distilled or deionised water to prevent build-up of salts.

#### **Stabilisation**

25. Once the spiked sediment with overlying water has been prepared, it is desirable to allow partitioning of the test chemical from sediment to the aqueous phase (3)(4)(6)(13). This should preferably be done under the conditions of temperature and aeration used in the test. Appropriate equilibration time is sediment and chemical specific. Organic chemicals with  $\log KOW < 4$  may reach equilibrium in 1 to 2 weeks, while those with  $\log KOW > 6$  can take 2 months or more (14). Metals may reach equilibrium within weeks to months, but the times differ depending on metals and sediments (14). As this would leave time for degradation of many chemicals, equilibrium is not required but an stabilisation period of 48 hours is recommended. An extended period for stabilisation or even equilibration should be considered and justified on a case-specific basis. The concentration of the test chemical should be measured in the overlying water, the pore water and the sediment, at least at the highest concentration and a lower one (see paragraph 38). These analytical determinations of the test chemical allow for calculation of mass balance (25) and expression of results based on measured concentrations (e.g. 26).

#### **Addition of test organisms**

26. Four to five days before adding the test organisms to the test vessels, egg masses should be taken from the cultures and placed in small vessels in culture medium. Aged medium from the stock culture or freshly prepared medium may be used. If the latter is used, a small amount of food, for example green algae and/or a few droplets of filtrate from a finely ground suspension of flaked fish food should be added to the culture medium (see Annex 2). Only freshly laid egg masses should be used. Normally, the larvae begin to hatch a couple of days after the eggs are laid (2 to 3 days for *C. riparius* at 20 °C and 1 to 4 days for *C. dilutus* at 23 °C and *C. yoshimatsui* at 25 °C) and larval growth occurs in four instars, each of 4 to 8 days duration. First-instar larvae (2-3 days post-hatch for *C. riparius*; 1-4 days post-hatch for *C. dilutus* and *C. yoshimatsui*) should be used in the test. In case of uncertainties on the instar of midges, it can possibly be checked using head capsule width (6).

27. Twenty first-instar larvae are allocated randomly to each test vessel containing the spiked sediment and water, using a blunt pipette. Gentle aeration of the water has to be stopped while adding the larvae to

test vessels and remain so for another e.g. 4 hours after addition of larvae (see paragraphs 25 and 32). According to the main test design used (see paragraphs 20 and 21), the number of larvae used per concentration is at least 60 for the EC point estimation and 80 for determination of NOEC.

### **Test concentrations**

28. A range-finding test may be helpful to determine the range of concentrations for the definitive test. For this purpose, a series of widely spaced concentrations of the test chemical are used. The chironomids are exposed to each concentration of the test chemical for a period which allows estimation of appropriate test concentrations, and no replicates are required. The larval density per vessel should be the same as for the definitive test. For the definitive test, at least five concentrations should be used and selected as described in paragraphs 19 to 21.

### **Controls**

29. Control vessels without any test chemical but including sediment should be included in the test with the appropriate number of replicates (see paragraphs 20 - 21). If a solvent has been used for application of test chemical (see paragraph 17), a solvent control (sediment spiked with solvent only) should be performed in addition to the usual control (or negative control). It should correspond to the highest solvent concentration of solvent used in the test treatments and generally not exceed 0.01%.

### **Test system**

30. Static systems are used. If evaporation occurs during the test, the volume of overlying water should be regularly adjusted by adding distilled or deionised water. Semi-static or flow-through systems with intermittent or continuous renewal of overlying water might be used in exceptional cases as for instance if water quality specifications become inappropriate for the test organism (e.g. dissolved oxygen levels or concentration of excretory products exceed the respective appropriate limits, or minerals leach from sediment and affect pH and/or water hardness). However, other methods for ameliorating the quality of overlying water, such as aeration, will normally suffice and be preferable.

### **Food**

31. It is necessary to feed the larvae, preferably daily or at least three times per week. Fish-food (a suspension in water or finely ground food, for example TetraMin® or Tetra-Phyll®; see details in Annex 2) in the amount of 0.25-0.5 mg (0.35-0.5 mg for *C. yoshimatsui*) per larvae per day seems adequate for young larvae for the first 10 days. Slightly more food may be necessary for older larvae: 0.5-1 mg per larvae per day should be sufficient for the rest of the test. Midges do not feed when they reach the pupal stage, so the feeding regime may be adapted to the number of larvae that have pupated and/or emerged to avoid low dissolved oxygen levels. The food ration should be reduced in all treatments and control if fungal growth is seen or if mortality is observed in controls. If fungal development cannot be managed the test is to be repeated. When testing strongly adsorbing chemicals (e.g. with log Kow > 5), or chemicals covalently binding to sediment, the amount of food necessary to ensure survival and natural growth of the organisms may be added to the formulated sediment before application of the test chemical and subsequent stabilisation (or even equilibration). For this, plant material should be used instead of fish food, for example the addition of 0.5 % (dry weight) finely ground leaves of stinging nettle (*Urtica dioeica*), mulberry (*Morus alba*), white clover (*Trifolium repens*), spinach (*Spinacia oleracea*) or of other plant material (dried and milled wheat grass or alpha-cellulose) may be used (27). Alternatively, field-collected sediment with a sufficiently high nutritional content for the total test period could be used instead of standard OECD sediment as far as chironomids



survive in it for the duration of the testing without showing signs of stress (e.g. emigration of larvae towards water column, unusual behaviour).

### **Incubation conditions**

32. Gentle aeration of the overlying water in test vessels is supplied preferably e.g. 4 hours after addition of the larvae and is continued throughout the test (care should be taken that dissolved oxygen concentration does not fall below 60 % of ASV). Aeration is provided through a glass Pasteur pipette fixed 2 - 3 cm above the sediment layer (i.e. one or few bubbles/sec). When testing volatile chemicals, consideration may be given not to aerate the sediment-water system.

33. The test is conducted at a constant temperature of 20 °C ( $\pm 2$  °C) for *C. riparius*. For *C. dilutus* and *C. yoshimatsui*, recommended temperatures are 23 °C and 25 °C ( $\pm 2$  °C), respectively. If the test is conducted in an isothermal room, the room temperature should be verified at appropriate time intervals (at least once daily). A 16 hours photoperiod is used and the light intensity should be 500 to 1000 lx.

### **Exposure duration**

34. The exposure commences with the addition of larvae to the spiked and control vessels. The maximum exposure duration is 28 days for *C. riparius* and *C. yoshimatsui*, and 65 days for *C. dilutus*. If midges emerge earlier, the test can be terminated after a minimum of five days after emergence of the last adult in the control.

## **Observations**

### **Emergence**

35. The number of fully emerged males, the number of fully emerged females as well as the total number of emerged midges (males and females) are recorded daily. Males are easily identified by their plumose antennae. The respective development rates are calculated according to paragraphs 52 and 53.

36. The test vessels should be observed at least three times per week to make visual assessment of any abnormal behaviour (for example leaving sediment, unusual swimming), compared with the control. During the period of expected emergence, a daily count of emerged midges is necessary. The sex and number of fully emerged midges are recorded daily. After identification the midges are removed from the vessels. Any egg masses deposited prior to the termination of the test should be recorded and then removed to prevent re-introduction of larvae into the sediment. The number of visible pupae that have failed to emerge is also recorded.

### **Growth and survival**

37. If data on 10-day larval survival and growth are to be provided, additional test vessels should be included at the start, so that they may be used subsequently. The sediment from these additional vessels is sieved using a 250  $\mu\text{m}$  sieve to retain the larvae. Criteria for death are immobility or lack of reaction to a mechanical stimulus. Larvae not recovered should also be counted as dead (larvae which have died at the beginning of the test may have been degraded by microbes). The (ash-free) dry weight of the surviving larvae per test vessel is determined and the mean individual dry weight per vessel calculated. It is useful to

determine which instar the surviving larvae belong to; for that measurement of the width of the head capsule of each individual can be used.

## Analytical measurements

### ***Concentration of the test chemical***

38. For the analytical determination of the test chemical concentration, samples of the overlying water, the pore water and the sediment should be taken for analysis at least at the start (i.e. shortly before addition of larvae) (see paragraph 25) and at the end of the test, at least at the highest concentration and a lower one, but preferably at all tested concentrations and from at least one vessel per treatment. These determinations of test chemical concentration inform about the behaviour/ partitioning of the tested chemical in the water-sediment system.

39. If the analysis needs large samples which cannot be taken from test vessels without influencing the test system (e.g. because samples are taken at different times of the test), analytical determinations should be performed on samples from additional test vessels treated in the same way (including the presence of test organisms) but not used for biological observations.

40. Centrifugation at, for example, 10,000 g and 4 °C for 30 min. is the recommended procedure to isolate pore water. However, if the test chemical is demonstrated not to adsorb to filters, filtration may also be acceptable. In some cases, it might not be possible to analyse concentrations in the pore water as the sample size is too small.

### ***Physical-chemical parameters***

41. The pH, dissolved oxygen and temperature should be measured at least in one vessel of each concentration level and one vessel of control(s) at the start of exposure and once per week 2 and 3 of exposure, and in all vessels designated for determination of biological variables at the end of the test period. Hardness and ammonia should be measured in the controls and one test vessel at the highest concentration at the start and the end of the test.

## DATA AND REPORTING

### Treatment of results

42. The purpose of this test is to determine the effect of the test chemical on the development rate, the number of fully emerged males, the number of fully emerged females as well as the total number of fully emerged midges, or in the case of the 10-day test effects on survival and weight of the larvae. If there are no indications of statistically different sensitivities of sexes, male and female results may be pooled for statistical analyses. The sensitivity differences between sexes can be statistically judged by, for example, a  $\chi^2$ -r  $\times$  2 table test. Larval survival and mean individual dry weight per vessel should be determined after 10 days where required.

43. Effect concentrations should always be reported in terms of mg test chemical/kg dry sediment. It is recommended to report mass balance calculations (25) (see Annex 5) for gaining information on partitioning of the chemical in the various compartments. This is optional depending on the requirements in the respective regulations; it is also particularly important for chemicals that are difficult to test (concentrations poorly

maintained in the test system). The endpoints can be also expressed in mg chemical/l overlying water when deemed relevant.

44. To express the endpoints as nominal concentrations based on dry sediment, there should be evidence that the concentration of the chemical being tested has been satisfactorily maintained in the sediment (i.e. between 80 % and 120 % of the nominal concentration throughout the test). If the concentrations are expected to deviate by more than 20 %, all test concentrations should be measured and more frequent analyses may be carried out.

For the sediment compartment, the endpoints should then be expressed in terms of geometric mean measured concentrations or in time-weighted arithmetic mean measured concentrations when the samples were taken in two or more unequal time intervals (with calculations considering also the additional intermediate analytical measurements) in accordance with recommendations in Annex 2 of OECD Guidance Document on Aquatic Toxicity Testing of Difficult Test Chemicals (12) and examples in Annex 6 of OECD Guideline 211: *Daphnia magna* Reproduction Test (26).

For the overlying water, the endpoints should be expressed similarly in terms of geometric mean measured concentrations or time-weighted arithmetic mean measured concentrations.

45. To convert concentration data expressed e.g. as mg/kg dry sediment into mg/kg organic carbon (OC), the dry sediment concentration should be divided by the percent TOC (expressed as decimal), as follows:

$$mg/kg\ OC = \frac{mg/kg\ dry\ sediment}{kg\ OC/kg\ dry\ sediment}$$

46. To compute a point estimate for the EC<sub>10</sub> and EC<sub>50</sub> or any other EC<sub>x</sub> values, the per-vessel statistics may be used as true replicates. In calculating a confidence interval for any EC<sub>x</sub> the variability among vessels should be taken into account, or it should be shown that this variability is so small that it can be ignored. When the model is fitted by Least Squares, a transformation should be applied to the per-vessel statistics in order to improve the homogeneity of variance. However, EC<sub>x</sub> values should be calculated after the response is transformed back to the original value. Further information on statistics are given in OECD Document 54 on the Current Approaches in the Statistical Analysis of Ecotoxicity Data: A Guidance to Application (28).

47. When the statistical analysis aims at determining the NOEC/LOEC by hypothesis testing, the variability among vessels needs to be taken into account, for example by a nested ANOVA. Alternatively, more robust tests (29) can be appropriate in situations where there are violations of the usual ANOVA assumptions.

### ***Emergence ratio***

48. Emergence ratios are quantal data, and can be analysed by the Cochran-Armitage test applied in step-down manner where a monotonic dose-response is expected and these data are consistent with this expectation. If not, a Fisher's exact or Mantel-Haentz test with Bonferroni-Holm adjusted p-values can be used. If there is evidence of greater variability between replicates within the same concentration than a binomial distribution would indicate (often referenced as "extra-binomial" variation), then a robust Cochran-Armitage or Fisher exact test such as proposed in (30), should be used. The CPFISH test (closure principle and Fisher-Freeman-Halton test (24)) may also be used for evaluation of emergence data (see paragraph 21).

The sum of midges emerged per vessel,  $n_e$ , is determined and divided by the number of larvae introduced,  $n_a$ :

$$ER = \frac{n_e}{n_a}$$

where:

- $ER$  = emergence ratio
- $n_e$  = number of midges emerged per vessel
- $n_a$  = number of larvae introduced per vessel

49. An alternative that is most appropriate for large sample sizes, when there is extra binomial variance, is to treat the emergence ratio as a continuous response and use procedures such as William's test when a monotonic dose-response is expected and is consistent with these ER data. Dunnett's test would be appropriate where monotonicity does not hold. A large sample size is defined here as the number emerged and the number not emerging both exceeding five, on a per replicate (vessel) basis.

50. To apply ANOVA methods, values of ER should first be transformed by the arcsin-sqrt transformation or Tukey-Freeman transformation to obtain an approximate normal distribution and to equalise variances. The Cochran-Armitage, Fisher's exact (Bonferroni), or Mantel-Haentzel tests can be applied when using the absolute frequencies. The arcsin-sqrt transformation is applied by taking the inverse sine (sine-1) of the square root of ER.

51. For emergence ratios, ECx-values are calculated using regression analysis (e.g. probit (30), logit, Weibull, appropriate commercial software etc.). If regression analysis fails (e.g. when there are less than two partial responses), other non-parametric methods such as moving average or simple interpolation are used.

### **Development rate**

52. The mean development time represents the mean time span between the introduction of larvae (day 0 of the test) and the emergence of the experimental cohort of midges. (For the calculation of the true development time, the age of larvae at the time of introduction should be considered). The development rate is the reciprocal of the development time (unit: 1/day) and represents that portion of larval development which takes place per day. The development rate is preferred for the evaluation of these sediment toxicity studies as its variance is lower, and it is more homogeneous and closer to normal distribution as compared to development time. Hence, powerful parametric test procedures may be used with development rate rather than with development time. For development rate as a continuous response, ECx-values can be estimated by using regression analysis (e.g. (31), (32)).

53. For the following statistical tests, the number of midges observed on inspection day  $x$  are assumed to be emerged at the mean of the time interval between day  $x$  and day  $x_l$  ( $l$  = length of the inspection interval, usually 1 day). The mean development rate per vessel is calculated according to:

$$\bar{x} = \sum_{i=1}^m \frac{f_i x_i}{n_e}$$

where:

- $\bar{x}$  : mean development rate per vessel
- $i$  : index of inspection interval
- $m$  : maximum number of inspection intervals
- $f_j$  : number of midges emerged in the inspection interval  $i$
- $n_e$  : total number of midges emerged at the end of experiment ( $= \sum f_j$ )
- $x_j$  : development rate of the midges emerged in interval  $i$

$$x_i = 1 / \left( \text{day}_i - \frac{1}{2} l_i \right)$$

where:

- $\text{day}_i$  : inspection day (days since application)
- $l_i$  : length of inspection interval  $i$  (days, usually 1 day)

## Test report

54. The test report should at least provide the following information:

### **Test chemical:**

- physical nature and, where relevant, physical-chemical properties (water solubility, vapour pressure, octanol water partition coefficient, partition coefficient in soil (or in sediment if available), stability in water, etc.);
- chemical identification data (common name, chemical name, structural formula, CAS number, etc.) including purity and analytical method for quantification of test chemical.

### **Test species:**

- test animals used: species, scientific name, source of organisms and breeding conditions;
- information on handling of egg masses and larvae;
- age of test animals when inserted into test vessels.

### **Test conditions:**

- sediment used, i.e. natural or formulated sediment;
- for natural sediment, location and description of sediment sampling site, including, if possible, contamination history; characteristics: pH, organic carbon content, C/N ratio and granulometry (if appropriate).

- preparation of the formulated sediment: ingredients and characteristics (organic carbon content, pH, moisture, etc. at the start of the test);
- preparation of the test water (if reconstituted water is used) and characteristics (oxygen concentration, pH, conductivity, hardness, etc. at the start of the test);
- depth of sediment and overlying water;
- volume of overlying and pore water; weight of wet sediment with and without pore water;
- test vessels (material and size);
- method of spiking sediment: test concentrations used, number of replicates and use of solvent if any;
- stabilisation phase of the spiked sediment-water system: duration and conditions, justification of extended period for stabilisation or even equilibration (if appropriate);
- incubation conditions: temperature, light cycle and intensity, aeration (frequency and intensity);
- detailed information on feeding including type of food, preparation, amount and feeding regime.

**Results:**

- the nominal test concentrations, the measured test concentrations and the results of all analyses to determine the concentration of the test chemical in the test vessel, including assessment of mass balance if appropriate;
- water quality within the test vessels, i.e. pH, temperature, dissolved oxygen, hardness and ammonia;
- replacement of evaporated test water, if any;
- number of emerged male midges, number of emerged female midges per vessel and per day;
- number of larvae which failed to emerge as midges per vessel;
- mean individual dry weight of larvae per vessel, and per instar, if appropriate;
- percent emergence per replicate and test concentration (male and female midges pooled);
- mean development rate of fully emerged midges per replicate and treatment rate (male and female midges pooled);
- estimates of toxic endpoints, for example EC<sub>x</sub> (and associated confidence intervals), NOEC and/or LOEC, and the statistical methods used for their determination;
- discussion of the results, including any influence on the outcome of the test resulting from deviations from this Guideline.

## LITERATURE

1. BBA (1995). Long-term toxicity test with *Chironomus riparius*: Development and validation of a new test system. Edited by M. Streloke and H.Köpp. Berlin 1995.
2. R. Fleming *et al.* (1994). Sediment Toxicity Tests for Poorly Water-Soluble Substances. Final Report to them European Commission. Report No: EC 3738. August 1994. WRc, UK.
3. SETAC (1993). Guidance Document on Sediment toxicity Tests and Bioassays for Freshwater and Marine Environments. From the WOSTA Workshop held in the Netherlands.
4. ASTM International/E1706-00 (2002). Test Method for Measuring the Toxicity of Sediment Associated Contaminants with Freshwater Invertebrates. pp 1125-1241. In ASTM International 2002 Annual Book of Standards. Volume 11.05. Biological Effects and Environmental Fate; Biotechnology; Pesticides. ASTM. International, West Conshohocken, PA.
5. Environment Canada (1997). Test for Growth and Survival in Sediment using Larvae of Freshwater Midges (*Chironomus tentans* or *Chironomus riparius*). Biological Test Method. Report SPE 1/RM/32. December 1997.
6. US-EPA (2000). Methods for Measuring the Toxicity and Bioaccumulation of Sediment associated Contaminants with Freshwater Invertebrates. Second edition. EPA 600/R-99/064. March 2000. Revision to the first edition dated June 1994.
7. US-EPA/OPPTS 850.1735. (1996). Whole Sediment Acute Toxicity Invertebrates.
8. US-EPA/OPPTS 850.1790. (1996). Chironomid Sediment toxicity Test.
9. Milani, D., K.E. Day, D.J. McLeay, and R.S. Kirby. (1996). Recent intra- and inter-laboratory studies related to the development and standardisation of Environment Canada's biological test methods for measuring sediment toxicity using freshwater amphipods (*Hyaella azteca*) and midge larvae (*Chironomus riparius*). Technical Report. Environment Canada. National Water Research Institute. Burlington, Ontario, Canada.
10. Sugaya, Y. (1997). Intra-specific variations of the susceptibility of insecticides in *Chironomus yoshimatsui*. Jp. J. Sanit. Zool. 48 (4): 345-350.
11. Kawai, K. (1986). Fundamental studies on Chironomid allergy. I. Culture methods of some Japanese Chironomids (Chironomidae, Diptera). Jp. J. Sanit. Zool. 37(1):47-57.
12. OECD (2019), Guidance Document on Aquatic Toxicity Testing of Difficult Test Chemicals, Environmental Health and Safety Publications, Series on Testing and Assessment. No. 23 (Second Edition), [ENV/JM/MONO\(2000\)6/REV1](#), OECD, Paris.
13. Environment Canada. (1995). Guidance Document on Measurement of Toxicity Test Precision Using Control Sediments Spiked with a Reference Toxicant. Report EPS 1/RM/30. September 1995.
14. ASTM E1706-20 (2020). Standard Test Method for Measuring the Toxicity of Sediment-Associated Contaminants with Freshwater Invertebrates.
15. OECD (2016), Test No. 222: Earthworm Reproduction Test (*Eisenia fetida/ Eisenia andrei*). OECD Guidelines for the Testing of Chemicals, Section 2, OECD Publishing, Paris, <https://doi.org/10.1787/9789264264496-en>
16. Suedel, B.C. and J.H. Rodgers. (1994). Development of formulated reference sediments for freshwater and estuarine sediment testing. Environ. Toxicol. Chem. 13: 1163-1175.
17. Naylor, C. and C. Rodrigues. (1995). Development of a test method for *Chironomus riparius* using a formulated sediment. Chemosphere 31: 3291-3303.

18. ECHA 2017. Guidance on the Biocidal Regulation. Volume IV Environment – Assessment and Evaluation (Parts B + C). Helsinki (FI).
19. ECHA 2017. Guidance on information requirements and chemical safety assessment. Chapter R.7b: endpoint specific guidance. Helsinki (FI).
20. Dunnett, C.W. (1964). A multiple comparisons procedure for comparing several treatments with a control. *J. Amer. Statis. Assoc.*, 50: 1096-1121.
21. Dunnett, C. W. (1964). New tables for multiple comparisons with a control. *Biometrics* 20: 482-491.
22. Williams, D. A. (1971). A test for differences between treatment means when several dose levels are compared with a zero dose control. *Biometrics* 27: 103-117.
23. Williams, D.A. (1972). The comparison of several dose levels with a zero dose control. *Biometrics* 28: 510-531.
24. Lehmann, R., J. Bachmann, B. Karaoglan, J. Lacker, C. Polleichtner, H.T. Ratte and M. Ratte (2018). An alternative approach to overcome shortcomings with multiple testing of binary data in ecotoxicology. *Stoch. Environ. Res. Risk. Assess.* 32: 213–222.
25. EFSA (2019). Pesticide Peer Review meeting on general recurring issues in ecotoxicology (EFSA Supporting publication 2019:EN-1673)
26. OECD (2012), Test No. 211: *Daphnia magna* Reproduction Test. OECD Guidelines for the Testing of Chemicals, Section 2, OECD Publishing, Paris, <https://doi.org/10.1787/9789264185203-en>
27. Egeler, P., K.S. Henry and C. Riedhammer (2010). Potential effects of food addition to sediment on test conditions in sediment toxicity tests. *J. Soils Sediments* 10: 377–388.
28. OECD (2006). Current approaches in the statistical analysis of ecotoxicity data: a guidance to application. Series on Testing and Assessment No. 54, OECD Paris. <https://doi.org/10.1787/9789264085275-en>.
29. Rao, J.N.K. and A.J. Scott. (1992). A simple method for the analysis of clustered binary data. *Biometrics* 48: 577-585.
30. Christensen, E.R. (1984). Dose-response functions in aquatic toxicity testing and the Weibull model. *Water Research* 18: 213-221.
31. Bruce, R.D. and D.J. Versteeg (1992). A statistical procedure for modelling continuous toxicity data. *Environ. Toxicol. Chem.* 11: 1485-1494.
32. Slob, W. (2002). Dose-response modelling of continuous endpoints. *Toxicol. Sci.* 66: 298-312.



## ANNEX 1 - DEFINITIONS

For the purpose of this guideline the following definitions are used:

- The conditioning period is used to stabilise the microbial component of the sediment and to remove e.g. ammonia originating from sediment components; it takes place prior to spiking of the sediment with the test chemical.
- The equilibration period is the time needed for partitioning of the test chemical to reach equilibrium from sediment to the aqueous phase.
- The stabilisation period is the time used after spiking to allow partitioning of the test chemical between the solid phase, the pore water and the overlying water; it takes place after preparation of the spiked sediment – water system and ideally under exposure conditions (e.g. temperature and aeration).
- Formulated sediment or reconstituted, artificial or synthetic sediment, is a mixture of materials used to mimic the physical components of a natural sediment.
- Growth: increase in dry weight of test organisms during the experiment and expressed as mean dry weight per surviving chironomid.
- Overlying water is the water placed over sediment in the test vessel.
- Pore water is the water occupying space between sediment and soil particles.
- Spiked sediment is sediment to which test chemical has been added.

## ANNEX 2 - RECOMMENDATIONS FOR CULTURE OF *CHIRONOMUS RIPARIUS*

1. *Chironomus* larvae may be reared in crystallising dishes or larger containers. Fine quartz sand is spread in a thin layer of about 5 to 10 mm deep over the bottom of the container. Kieselgur (e.g. Merck, Art 8117) has also been shown to be a suitable substrate (a thinner layer of up to a very few mm is sufficient). Suitable water is then added to a depth of several cm. Water levels should be topped up as necessary to replace evaporative loss, and prevent desiccation. Water can be replaced if necessary. Gentle aeration should be provided. The larval rearing vessels should be held in a suitable cage which will prevent escape of the emerging adults. The cage should be sufficiently large to allow swarming of emerged adults, otherwise copulation may not occur (minimum is ca. 30 x 30 x 30 cm).

2. Cages should be held at room temperature or in a constant environment room at  $20 \pm 2$  °C with a photo period of 16-hour light (intensity ca. 1000 lx), 8 hours dark. It has been reported that air humidity of less than 60 % RH can impede reproduction.

### Dilution water

3. Any suitable natural or synthetic water may be used. Well water, dechlorinated tap water and artificial media (e.g. Elendt "M4" or "M7" medium, see below) are commonly used. The water has to be aerated before use. If necessary, the culture water may be renewed by pouring or siphoning the used water from culture vessels carefully without destroying the tubes of larvae.

### Feeding larvae

4. *Chironomus* larvae should be fed with a fish flake food (TetraMin<sup>®</sup>, TetraPhyll<sup>®</sup> or other similar brand of proprietary fish food), at approximately 250 mg per vessel per day. This can be given as a dry ground powder or as a suspension in water: 1.0 g of flake food is added to 20 ml of dilution water and blended to give a homogenous mix. This preparation may be fed at a rate of about 5 ml per vessel per day (shake before use). Older larvae may receive more.

5. Feeding is adjusted according to the water quality. If the culture medium becomes 'cloudy', the feeding should be reduced. Food additions must be carefully monitored. Too little food will cause emigration of the larvae towards the water column, and too much food will cause increased microbial activity and reduced oxygen concentrations. Both conditions can result in reduced growth rates.

6. Some green algae (e.g., *Scenedesmus subspicatus*, *Chlorella vulgaris*) cells may also be added when new culture vessels are set up.

### Feeding emerged adults

7. Some experimenters have suggested that a cotton wool pad soaked in a saturated sucrose solution may serve as a food for emerged adults.

## Emergence

8. At  $20 \pm 2$  °C adults will begin to emerge from the larval rearing vessels after approximately 13 - 15 days. Males are easily distinguished by having plumose antennae.

## Egg masses

9. Once adults are present within the breeding cage, all larval rearing vessels should be checked three times weekly for deposition of the gelatinous egg masses. If present, the egg masses should be carefully removed. They should be transferred to a small dish containing a sample of the breeding water. Egg masses are used to start a new culture vessel (e.g. 2 - 4 egg masses / vessel) or are used for toxicity tests.

10. First instar larvae should hatch after 2 - 3 days.

## Set-up of new culture vessels

11. Once cultures are established it should be possible to set up a fresh larval culture vessel weekly or less frequently depending on testing requirements, removing the older vessels after adult midges have emerged. Using this system a regular supply of adults will be produced with a minimum of management.

## Preparation of test solutions “M4” and “M7”

12. Elendt (1990) has described the "M4" medium. The "M7" medium is prepared as the "M4" medium except for the substances indicated in Table 1, for which concentrations are four times lower in "M7" than in "M4". The test solution should not be prepared according to Elendt and Bias (1990) for the concentrations of  $\text{NaSiO}_3 \cdot 5 \text{H}_2\text{O}$ ,  $\text{NaNO}_3$ ,  $\text{KH}_2\text{PO}_4$  and  $\text{K}_2\text{HPO}_4$  given for the preparation of the stock solutions are not adequate.

## Preparation of the “M7” medium

13. Each stock solution (I) is prepared individually and a combined stock solution (II) is prepared from these stock solutions (I) (see Table 1). Fifty ml from the combined stock Solution (II) and the amounts of each macro nutrient stock solution which are given in Table 2 are made up to 1 litre of deionised water to prepare the "M7" medium. A vitamin stock solution is prepared by adding three vitamins to deionised water as indicated in Table 3, and 0.1 ml of the combined vitamin stock solution are added to the final "M7" medium shortly before use. (The vitamin stock solution is stored frozen in small aliquots). The medium is aerated and stabilised.

Table 1: Stock solutions of trace elements for medium M4 and M7

Stock solutions (I)	Amount (mg) made up to 1 litre of deionised water	To prepare the combined stock solution (II): mix the following amounts (ml) of stock solutions (I) and make up to 1 litre of deionised water		Final concentrations in test solutions (mg/l)	
		M4	M7	M4	M7
H <sub>3</sub> BO <sub>3</sub> <sup>(1)</sup>	57190	1.0	0.25	2.86	0.715
MnCl <sub>2</sub> 4 H <sub>2</sub> O <sup>(1)</sup>	7210	1.0	0.25	0.361	0.090
LiCl <sup>(1)</sup>	6120	1.0	0.25	0.306	0.077
RbCl <sup>(1)</sup>	1420	1.0	0.25	0.071	0.018
SrCl <sub>2</sub> 6 H <sub>2</sub> O <sup>(1)</sup>	3040	1.0	0.25	0.152	0.038
NaBr <sup>(1)</sup>	320	1.0	0.25	0.016	0.004
Na <sub>2</sub> MoO <sub>4</sub> 2 H <sub>2</sub> O <sup>(1)</sup>	1260	1.0	0.25	0.063	0.016
CuCl <sub>2</sub> 2 H <sub>2</sub> O <sup>(1)</sup>	335	1.0	0.25	0.017	0.004
ZnCl <sub>2</sub>	260	1.0	1.0	0.013	0.013
CoCl <sub>2</sub> 6 H <sub>2</sub> O	200	1.0	1.0	0.010	0.010
KI	65	1.0	1.0	0.0033	0.0033
Na <sub>2</sub> SeO <sub>3</sub>	43.8	1.0	1.0	0.0022	0.0022
NH <sub>4</sub> VO <sub>3</sub>	11.5	1.0	1.0	0.00058	0.00058
Na <sub>2</sub> EDTA 2 H <sub>2</sub> O <sup>(1)(2)</sup>	5000	20.0	5.0	2.5	0.625
FeSO <sub>4</sub> 7 H <sub>2</sub> O <sup>(1)(2)</sup>	1991	20.0	5.0	1.0	0.249

1. These substances differ in M4 and M7, as indicated above.
2. These solutions are prepared individually, then poured together and autoclaved immediately.

Table 2: Macro nutrient stock solutions for medium M4 and M7

	Amount made up to 1 litre of deionised water (mg)	Amount of macro nutrient stock solutions added to prepare medium M4 and M7 (ml/l)	Final concentrations in test solutions M4 and M7 (mg/l)
CaCl <sub>2</sub> 2 H <sub>2</sub> O	293800	1.0	293.8
MgSO <sub>4</sub> 7 H <sub>2</sub> O	246600	0.5	123.3
KCl	58000	0.1	5.8
NaHCO <sub>3</sub>	64800	1.0	64.8
NaSiO <sub>3</sub> 9 H <sub>2</sub> O	50000	0.2	10.0
NaNO <sub>3</sub>	2740	0.1	0.274
KH <sub>2</sub> PO <sub>4</sub>	1430	0.1	0.143
K <sub>2</sub> HPO <sub>4</sub>	1840	0.1	0.184

**Table 3: Vitamin stock solutions for medium M4 and M7**

All three vitamin solutions are combined to make a single vitamin stock solution.

	Amount made up to 1 litre of deionised water (mg)	Amount of vitamin stock solution added to prepare medium M4 and M7 (ml/l)	Final concentrations in test solutions M4 and M7 (mg/l)
Thiamine hydrochloride	750	0.1	0.075
Cyanocobalamin (B12)	10	0.1	0.0010
Biotine	7.5	0.1	0.00075

## References

- Elendt, B.P. (1990): Selenium Deficiency in Crustacean. 154, 25-33
- Elendt, B.P. & W.-R. Bias (1990): Trace Nutrient Deficiency in *Daphnia magna* Cultured in Standard Medium for Toxicity Testing. Effects on the Optimization of Culture Conditions on Life History Parameters of *D. magna*. *Water Research* 24 (9), 1157-1167

# ANNEX 3 - PREPARATION OF FORMULATED SEDIMENT

## Sediment composition

The composition of the formulated sediment should be as follows:

Constituent	Characteristics	% of sediment dry weight
Peat	e.g. Sphagnum moss peat, as close to pH 5.5-6.0 as possible, no visible plant remains, finely ground (particle size $\leq$ 1 mm) and air dried	4-5
Quartz sand	Grain size: > 50% of the particles should be in the range of 50-200 $\mu$ m	75-76
Kaolinite clay	Kaolinite content 30%	20
Organic carbon	Adjusted by addition of peat and sand	2 ( $\pm$ 0.5)
Calcium carbonate	CaCO <sub>3</sub> , pulverised, chemically pure	0.05-1
Water	Conductivity 10 $\mu$ S/cm	30-50

## Preparation

1. The peat is air dried and ground to a fine powder. A suspension of the required amount of peat powder in deionised water is prepared using a high-performance homogenising device. The pH of this suspension is adjusted to  $5.5 \pm 0.5$  with CaCO<sub>3</sub>. The suspension is conditioned for at least two days with gentle stirring at  $20 \pm 2$  °C, to stabilise pH and establish a stable microbial component. pH is measured again and should be  $6.0 \pm 0.5$ . Then the peat suspension is mixed with the other constituents (sand and kaolin clay) and deionised water to obtain a homogeneous sediment with a water content in a range of 30-50 per cent of dry weight of the sediment. The pH of the final mixture is measured once again and is adjusted to 6.5 to 7.5 with CaCO<sub>3</sub> if necessary. Samples of the sediment are taken to determine the dry weight and the organic carbon content. Then, before it is used in the chironomid toxicity test, it is recommended that the formulated sediment be conditioned for seven days under the same conditions which prevail in the subsequent test.

## Storage

2. The dry constituents for preparation of the artificial sediment may be stored in a dry and cool place at room temperature. The formulated (wet) sediment should not be stored prior to its use in the test. It should be used immediately after the 7 days conditioning period that ends its preparation.

## References

- OECD (1984), Test No. 207: Earthworm, Acute Toxicity Tests, OECD Guidelines for the Testing of Chemicals, Section 2, OECD Publishing, Paris, <https://doi.org/10.1787/9789264070042-en>.
- Meller, M., P. Egeler, J. Rombke, H. Schallnass, R. Nagel and B. Streit. (1998). Short-term Toxicity of Lindane, Hexachlorobenzene and Copper Sulfate on Tubificid Sludgeworms (Oligochaeta) in Artificial Media. *Ecotox. and Environ. Safety*, 39, 10-20.

## ANNEX 4 - CHEMICAL CHARACTERISTICS OF AN ACCEPTABLE DILUTION WATER

SUBSTANCE	CONCENTRATIONS
Particulate matter	< 20 mg/l
Total organic carbon	< 2 mg/l
Unionised ammonia	< 1 µg/l
Hardness as CaCO <sub>3</sub>	< 400 mg/l*
Residual chlorine	< 10 µg/l
Total organophosphorus pesticides	< 50 ng/l **
Total organochlorine pesticides plus polychlorinated biphenyls	< 50 ng/l **
Total organic chlorine	< 25 ng/l **

\* However, it should be noted that if there is an interaction suspected between hardness ions and the test chemical, lower hardness water should be used (and thus, Elendt Medium M4 must not be used in this situation).

\*\*Please note that this item is only to be checked in case natural or tap water is used.



## ANNEX 5-MASS BALANCE CALCULATIONS FOR SEDIMENT-WATER TOXICITY TESTS; EXAMPLE FOR SPIKED SEDIMENT TEST PERFORMED WITH CHIRONOMUS SP.

1. The starting point for a mass balance calculation for a study with sediment-dweller organisms in sediment-water tests using spiked sediment (e.g. OECD TG 218) or spiked water (e.g. OECD TG 219) is the measured concentrations of the chemical in the test system considering the concentrations at the beginning and end of the study as well any additional intermediate concentrations available.
2. The mass balance calculation in the case of studies with sediment-dwellers with spiked sediment (OECD TG 218) is illustrated below for one test chemical unstable in sediment and one test chemical stable in sediment such as two pesticides with different characteristics.
3. The information on the analytical determination of the chemical needed for the mass balance calculation is given for the two examples in Table 1A and Table 2A. The measurements in these two cases were at three time points (1 hour, 7 days and 28 days). For simplicity, they are reported here only at the lowest and highest concentrations tested. However, measurements at additional tested concentrations are recommended, particularly in the case of the chemicals that are difficult to test (e.g. when concentrations are poorly maintained in the test system).
4. For the two examples below, the test system contains an amount of sediment of 0.1 kg and a volume of overlying water of 400 ml; it is assumed that the pore water has a volume of 10 ml. Based on this information, it is possible to calculate the amount of chemical that partitions to the various compartments by multiplying the concentration by the volume/amount of water/sediment and to calculate the percentage in each compartment with respect to that initially contained in the spiked sediment. In the example 1, in the absence of information about the pore water concentrations, only the amount of chemical in overlying water is considered.
5. In order to assess the fate of the test chemical in the sediment/water system, the mass balance calculation was performed for each time point for which measurements were available (see Tables 1B and 2B).

## Example 1: Test chemical unstable in sediment

**Table 1A:** Measured concentrations of the test chemical in different compartments of a spiked sediment study with *Chironomus riparius* performed in line with OECD TG 218.

Measured concentrations-(mg a.s./kg dry sediment or µg a.s./L)									
	day 0 (start of exposure)			7 days			28 days		
Nominal (mg a.s./kg)	Overlying water (µg a.s./l)	Pore water (µg a.s./l)	Sediment (mg a.s./kg)	Overlying water (µg a.s./l)	Pore water (µg a.s./l)	Sediment (mg a.s./kg)	Overlying water (µg a.s./l)	Pore water (µg a.s./l)	Sediment (mg a.s./kg)
Control	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ
0.125	0.00406	/	0.12	0.784	/	0.06	3.2	/	0.046
2	0.0660	/	1.96	3.39	/	1.1	40.1	/	0.81

a.s.: active substance (test chemical); LOQ: limit of quantification

**Table 1B:** Example of mass balance calculations for the test chemical. The amount in both compartments – sediment and water (overlying water)- is expressed in terms of total amounts (Table 1B-1) and in percentage (Table 1B-2) of test chemical with respect to the amount initially contained in the system (measured in water and sediment). Please note that the pore water concentrations are not given and thus implying significant uncertainties in the calculations.

Table 1B-1

Amounts (µg) calculated from measured concentrations									
	day 0 (start of exposure)			7 days			28 days		
Nominal (µg)	Overlying water (µg)*	Pore water (µg)*	Sediment (µg)*	Overlying water (µg)	Pore water (µg)	Sediment (µg)	Overlying water (µg)	Pore water	Sediment (µg)
Control	<LOQ	/	<LOQ	<LOQ	/	<LOQ	<LOQ	/	<LOQ
12.5	0.0016	/	12	0.3136	/	6.0	1.280	/	4.6
200	0.0264	/	196	1.356	/	110	16.04	/	81

Calculated considering \*100 g sediment, 400 ml overlying water

Table 1B-2

Mass balance									
	day 0 (start of exposure)			7 days			28 days		
Nominal (µg)	Overlying water (%)	Pore water (%)	Sediment (%)	Overlying water (%)	Pore water (%)	Sediment (%)	Overlying water (%)	Pore water (%)	Sediment (%)
Control	<LOQ	/	<LOQ	<LOQ	/	<LOQ	<LOQ	/	<LOQ
12.5	0.013	/	99.987	2.613	/	49.993	10.665	/	38.328
200	0.013	/	99.987	0.692	/	56.115	8.182	/	41.321

6. In the example 1, the test chemical is measured almost exclusively in the sediment in the first hour (i.e. at levels of ca. 99.9 %) which is expected since the sediment has been spiked prior to the start of the experiment. But at the end of the study (i.e. at 28 days), the test chemical is only present in the spiked sediment at levels up to ca. 40 % of that initially measured. In the case of the water compartment, only overlying water concentrations are available (thus implying some uncertainties in the calculations) with increasing concentrations over the test duration; this indicates remobilisation/ desorption of the a.s. from the sediment into the overlying water with a total amount at the end of the study reaching ca. 10 % of the amounts initially measured.

7. During this test, degradation occurred; the total amount of the test chemical in the test system in the study accounts for up to ca. 49 % of the total amount initially measured.

In such a case, it is needed to express the endpoints in terms of mean measured concentrations (preferably time-weighted arithmetic mean measured concentrations rather than geometric mean measured concentrations, as the time intervals are unequal) as the concentrations in sediment were not maintained within 20 % of the initial measured concentrations. It is also highly recommended to analyse all tested concentrations and any further additional intermediate measurements. It is additionally recommended that the key endpoints are presented in terms of mg of chemical per kg of dry sediment and mg of chemical per l of water. This would ensure that both exposure via water and sediment are covered for sediment-dwellers.

## Example 2: Test chemical stable in sediment

**Table 2A:** Measured concentrations of the test chemical in a spiked sediment study with *Chironomus riparius* performed in line with OECD TG 218.

Measured concentrations (mg a.s./kg dry sediment or µg a.s./L)									
	day 0 (start of exposure)			7 days			28 days		
Nominal (mg a.s./kg)	Overlying water (µg a.s./l)	Pore water (µg a.s./l)	Sediment (mg a.s./kg)	Overlying water (µg a.s./l)	Pore water (µg a.s./l)	Sediment (mg a.s./kg)	Overlying water (µg a.s./l)	Pore water (µg a.s./l)	Sediment (mg a.s./kg)
Control	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ

0.125	0.00406	0.798	0.12	0.00784	0.615	0.11	<LOQ	0.230	0.094
2	0.0660	25.5	1.96	0.0339	8.63	1.76	<LOQ	5.74	1.5

a.s.: active substance (test chemical); LOQ: limit of quantification

**Table 2B:** Example of mass balance calculations for a test chemical. The amount in both compartments – sediment and water (overlying and pore water)- is expressed in terms of total amounts (Table 2B-1) and in percentage (Table 2B-2) of test chemical with respect to the initially contained in the system (measured in water and sediment).

Table 2B-1

Amounts calculated from measured concentrations									
	day 0 (start of exposure)			7 days			28 days		
Nominal (µg in sediment)	Overlying water (µg)*	Pore water (µg)*	Sediment (µg)*	Overlying water (µg)	Pore water (µg)	Sediment (µg)	Overlying water (µg)	Pore water	Sediment (µg)
Control	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ
12.5	0.0016	0.008	12	0.0031	0.006	11	<LOQ	0.002	9.4
200	0.0264	0.255	196	0.0136	0.086	176	<LOQ	0.057	150

Calculated considering \*100 g sediment, 400 ml overlying water, and assuming 10 ml pore water (uncertainties in the calculations)

Table 2B-2

Mass balance									
	day 0 (start of exposure)			7 days			28 days		
Nominal (µg)	Overlying water (%)	Pore water (%)	Sediment (%)	Overlying water (%)	Pore water (%)	Sediment (%)	Overlying water (%)	Pore water (%)	Sediment (%)
Control	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ	<LOQ
12.5	0.013	0.067	99.920	0.026	0.050	91.593	/	0.019	78.271
200	0.013	0.130	99.857	0.007	0.044	89.667	/	0.029	76.421

8. In the example 2, the test chemical is measured almost exclusively in the sediment in the first hour (i.e. at levels of ca. 99.9 %), which is expected since the sediment has been spiked prior to the start of the experiment. At the end of the study, the test chemical remains present in the spiked sediment at levels up to ca. 80 % of that initially measured. In the case of the water compartments (overlying and pore waters),

the amounts are much lower than in the sediment. The total amount of the test chemical in the test system (sediment and pore water) in this example accounts for up to ca. 83 % of the amount initially measured in the sediment compartment.

In such a case, it is possible to express the key endpoints in terms of initial concentrations in sediment as the concentrations were maintained within 20 % of the initial measured concentrations (i.e. in terms of mg of chemical per kg of dry sediment) and as the exposure via water is not occurring. It is also recommended to analyse all tested concentrations and any further additional intermediate measurements.

Please note that the expression of endpoints in terms of mg of chemical per kg of dry sediment could also be complemented by mg of chemical per kg of organic carbon.